

- 46.** Waterham HR, Titorenko VI, Haima P, Cregg JM, Harder W and Veenhuis M. The *Hansenula polymorpha* *PER1* gene is essential for peroxisome biogenesis and encodes a peroxisomal matrix protein with both carboxy- and amino-terminal targeting signals. *J Cell Biol.* 1994; 127:737-749.
- 47.** Klei IJ, Harder W and Veenhuis M. Methanol metabolism in a peroxisome-deficient mutant of *Hansenula polymorpha*: a physiological study. *Arch Microbiol.* 1991; 156:15-23.
- 48.** Lück H. (1963). Catalase. (New York: Academic Press).
- 49.** Culotta VC, Klomp LWJ, Strain J, Casareno RLB, Krems B and Gitlin JD. The copper chaperone for superoxide dismutase. *J Biol Chem.* 1997; 272:23469-23472.
- 50.** Flohé L and Otting F. Superoxide dismutase assays. *Methods Enzymol.* 1984; 105:93-104.
- 51.** Verduyn C, Giuseppin MLF, Scheffers WA and van Dijken JP. Hydrogen peroxide metabolism in yeasts. *Appl Environ Microbiol.* 1988; 54: 2086-2090.
- 52.** Baerends RJ, Faber KN, Kram AM, Kiel JA, van der Klei IJ and Veenhuis M. A stretch of positively charged amino acids at the N terminus of *Hansenula polymorpha* Pex3p is involved in incorporation of the protein into the peroxisomal membrane. *J Biol Chem.* 2000; 275:9986-9995.
- 53.** Sudbery PE, Gleeson MA, Veale RA, Ledeboer AM and Zoetmulder MC. *Hansenula polymorpha* as a novel yeast system for the expression of heterologous genes. *Biochem Soc Trans.* 1988; 16:1081-1083.
- 54.** Nagotu S, Saraya R, Otzen M, Veenhuis M and van der Klei IJ. Peroxisome proliferation in *Hansenula polymorpha* requires Dnm1p which mediates fission but not de novo formation. *Biochim Biophys Acta.* 2008; 1783:760-769.
- 55.** Nagotu S, Krikken AM, Otzen M, Kiel JAKW, Veenhuis M and Van Der Klei IJ. Peroxisome fission in *Hansenula polymorpha* requires Mdv1 and Fis1, two proteins also involved in mitochondrial fission. *Traffic.* 2008; 9:1471-1484.
- 56.** Saraya R, Krikken AM, Veenhuis M and van der Klei IJ. Peroxisome reintroduction in *Hansenula polymorpha* requires Pex25 and Rho1. *J Cell Biol.* 2011; 193:885-900.

SUPPLEMENTARY FIGURES




Figure S1. Growth curves of WT and cat cells.
Cells were grown on (A) Glc/AS, (B) Glc/MA, (C) Gly/AS or (D) Gly/MeOH/AS. Optical densities are expressed as OD_{600 nm}. (E). Residual methanol concentrations in WT and cat cultures at different time points after shifting the cells to Gly/MeOH/AS. Bars indicate SD of 3 independent cultures.




Figure S2. Sequence alignment of Yap1 homologues from different yeast species. The gene bank accession numbers are: *H. polymorpha* (EFW96135), *P. pastoris* (XP_002494040), *S. pombe* (CAB66170), *S. cerevisiae* (NP_013707).




Figure S3. Growth curves of *yap1* and *cat yap1* cells. Cells were grown on (A) Glc/AS, (B) Glc/MA, (C) Gly/AS, (D) (Gly/MeOH/AS). Bars indicate SD of 2 independent cultures.