

Correction for: The role of serum growth hormone and insulin-like growth factor-1 in adult humans brain morphology

Taoyang Yuan¹, Jianyou Ying¹, Lu Jin², Chuzhong Li¹, Songbai Gui², Zhenye Li², Rui Wang¹, Zhentao Zuo^{3,4,5}, Yazhuo Zhang^{1,2,6}

¹Beijing Neurosurgical Institute, Capital Medical University, Beijing, China

²Department of Neurosurgery, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

³State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese Academy of Sciences, Beijing, China

⁴CAS Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, Beijing, China

⁵Sino-Danish College, University of Chinese Academy of Sciences, Beijing, China

⁶Beijing Institute for Brain Disorders, Brain Tumour Center, China National Clinical Research Center for Neurological Diseases, Key Laboratory of Central Nervous System Injury Research, Beijing, China

Correspondence to: Zhentao Zuo, Yazhuo Zhang; **email:** ztzuo@bcslab.ibp.ac.cn, zyz2004520@yeah.net

Original article: *Aging (Albany NY)* 2020; 12: pp 1377–1396

PMID: [31967977](https://pubmed.ncbi.nlm.nih.gov/31967977/)

PMCID: [PMC7053622](https://pubmed.ncbi.nlm.nih.gov/PMC7053622/)

doi: [10.18632/aging.102688](https://doi.org/10.18632/aging.102688)

This article has been corrected: The authors recently found that panels **H** and **J** in **Figure 2** are the same, as they accidentally placed normalized WM plot instead of normalized GM plot. They replaced panel **H** **Figure 2** with the correct plot from the original data. This alteration does not affect the results or conclusions of this work.

New **Figure 2** is presented below.




Figure 2. Correlation analysis between serum GH/IGF-1 levels and brain tissue volume in patients with GH-PA and patients with NonFun-PA groups. The normalized GM volume (nGMV) shows significant positive correlation with GH (A) and IGF-1 (B) in patients with GH-PA. The normalized WM volume (nWMV) shows no significant correlation with GH (C) or IGF-1 (D) in patients with GH-PA. The normalized CSF volume (nCSFV) shows significant negative correlation with GH (E) and IGF-1 (F) in patients with GH-PA. In patients with NonFun-PA, nGMV, nWMV, and nCSFV show no significant correlation with GH/IGF-1 (G–L).