Correction for: miR-320 accelerates chronic heart failure with cardiac fibrosis through activation of the IL6/STAT3 axis

Fang Li¹, Shan-Shan Li¹, Hui Chen¹, Jian-Zhi Zhao², Jie Hao¹, Jin-Ming Liu¹, Xiu-Guang Zu¹, Wei Cui³

¹Third Division, Department of Cardiology, The Second Hospital of Hebei Medical University, Shijiazhuang, Hebei 050011, PR China

²Department of Biochemistry and Molecular Biology, The Hebei Medical University, Shijiazhuang, Hebei 050011, PR China

³Department of Cardiology, The Second Hospital of Hebei Medical University and Hebei Institute of Cardiovascular Research, Shijiazhuang, Hebei 050011, PR China

Correspondence to: Wei Cui; **email:** <u>cuiwei@hb2h.com</u>, <u>https://orcid.org/0000-0002-1214-4146</u> **Keywords:** miR-320, chronic heart failure, cardiac fibrosis, IL6/STAT3/PTEN axis

Original article: Aging (Albany NY) 2021; 13: pp 22516-22527

PMID: <u>34582362</u> PMCID: <u>PMC8507257</u> doi: <u>10.18632/aging.203562</u>

This article has been corrected: The authors found an error in Figure 7A: the Western blot band for Collagen-I was mistakenly also placed at the position for IL-6. The authors prepared a new Figure 7 using images from the original experiments and recalculated the relative densities of these proteins. This correction does not impact the conclusions of the paper.

New **Figure 7** is presented below.



Figure 7. STAT3 was involved in miR-320 mimics-induced development of cardiac hypertrophy and fibrosis. (A) Western blots revealed that BP-1-102 significantly reversed the up-regulated STAT3, p-STAT3, type I and type III collagen, IL6, TGF- β , and down-regulated PTEN. (B) The quantitative analysis illustrated that inhibitor BP-1-102 exposure blunted the up-regulated type I and III collagen, IL6, TGF- β , p-STAT3, and STAT3 expression and enhanced the down-regulated PTEN expression in fibroblasts transfected with miR-320 mimic by BP-1-102. P < 0.05.